INDEX TO AUTHORS

BERENBLUM, I., and SCHOENTAL, R., carcinogenic constituents of coal-tar, 157

——— and SHUBIK, P., the role of croton oil applications, associated with a single painting of a carcinogen, in tumour induction of the mouse's skin, 379

——— ——— a new, quantitative, approach to the study of the stages of chemical carcinogenesis in the mouse's skin, 383

BIELSCHOWSKY, F., the effect of a diet containing yeast upon the development of tumours induced by 2-acetyl-amino-fluorene, 146

——— see *Orr, J. W.*

BURGEN, A. S. V., see *Fairburn, E. A.*

BUSK, T., see *Clemmesen, J.*

——— and CLEMMESEN, J., the frequencies of left- and right-sided breast cancer, 345

BUSSEY, H. J. R., see *Dukes, C. E.*

CALCUTT, G., and POWELL, A. K., the control of some factors involved in experimental epidermal carcinogenesis, 323.

——— ——— see *Weigert, F.*

CARR, J. G., production of mutations in mice by 1:2:5:6-dibenzanthracene, 152

CLEMMESEN, J., and BUSK, T., on the apparent increase in the incidence of lung cancer in Denmark, 1931–1945, 253

——— see *Busk, T.*

COWEN, P. N., some studies on the action of urethane on mice, 401

CRUICKSHANK, D. B., regional influences in cancer, 109

DMOCHOWSKI, L., see *Horning, E. S.*

DUKES, C. E., and BUSSEY, H. J. R., sarcoma and melanoma of the rectum, 30

DUSTIN, P., Jr., the cytological action of ethyl-carbamate (urethane) and other carbamic esters in normal and leukaemic mice, and in rabbits, 48

ELSON, L. A., GOULDEN, F., and WARREN, F. L., the absence of a direct relation between growth inhibition and sulphur metabolism after 1:2:5:6-dibenzanthracene, 80

——— and WARREN, F. L., the relation between growth inhibition, toxicity and protein metabolism in rats treated with 1:2:5:6-dibenzanthracene, 86

——— and HADDOW, A., the inhibitory action of 1:2:5:6-dibenzanthracene on the growth of the Walker carcinoma 256 in rats maintained on high and low protein diets, 97

ELSON L. A., and HARRIS, R. J. C., the influence of 1:2:5:6-dibenzanthracene on the nucleic acid content of the liver of rats maintained on high and low protein diets, 327

FAIRBURN, E. A., squamous cell carcinoma of the buccopharyngeal region, 20

——— and BURGEN, A. S. V., the skin lesions of monocytic leukaemia, 352

FOULDS, L., the induction of tumours in mice of the R3 strain by 2-acetylaminofluorene, 172

——— mammary tumours in hybrid mice: a sex-factor in transplantation, 362

GOULDEN, F., see *Elson, L. A.*

HADDOW, A., see *Elson, L. A.*

HANDLEY, R. S., and THACKRAY, A. C., invasion of the internal mammary lymph glands in carcinoma of the breast, 15

HARRIS, R. J. C., see *Elson, L. A.*

HOCH-LIGETI, C., effect of feeding 7 OH-2-acetaminofluorene to albino rats, 391

HORNING, E. S., and DMOCHOWSKI, L., induction of prostate tumours in mice, 59

KENNAWAY, E. L., cancer of the penis and circumcision in relation to the incubation period of cancer, 335

——— and KENNAWAY, N. M., a further study of the incidence of cancer of the lung and larynx, 260

KENNAWAY, N. M., see *Kennaway, E. L.*

KIRBY, A. H. M., the combined action of 2-acetylaminofluorene and sex hormones in the Wister rat, 68

KOLLER, P. C., abnormal mitosis in tumours, 38

MANN, IDA, induction of an experimental tumour of the lens, 63

MITCHELL, J. S., applications of recent advances in nuclear physics to cancer research, 1

NEUBAUER, O., arsenical cancer: a review, 192

ORR, J. W., the induction of pulmonary adenomata in mice by urethane, 311

——— the histology and histogenesis of pulmonary adenomata of mice, 316

——— and BIELSCHOWSKY, F., the histology of pulmonary tumours of the rat induced by 2-acetylaminofluorene, 396

POWELL, A. K., see *Calcutt, G.*
——— see *Weigert, F.*
PULLINGER, B. D., forced activation and early detection of the milk-borne agent of mammary adenomas in mice, 177

RIESCO, A., on the bearing of time on the neoplastic action of small quantities of α-oestradiol in the endometrium of guinea-pigs, 166
ROBINSON, R. H. M., see *Selbie, F. R.*

SCHOENTAL, R., see *Berenblum, I.*
SELBIE, F. R., and ROBINSON, R. H. M., serial transmission of infectious papillomatosis in the domestic rabbit, 371
SHUBIK, P., see *Berenblum, I.*
SPICER, C. C., the estimation of tumour susceptibility in pure lines, 298
SYLVÉN, B., changes in cartilage produced by infiltrating carcinoma, 103

THACKRAY, A. C., see *Handley, R. S.*
TRUSCOTT, B. McN., carcinoma of the breast. An analysis of the symptoms, factors affecting prognosis, results of treatment and recurrences in 1211 cases treated at the Middlesex Hospital, 129

WARREN, F. L., see *Elson, L. A.*
WEIGERT, F., CALCUTT, G., and POWELL, A. K., the course of the metabolism of benzpyrene in the skin of the mouse, 405
WEIL-MALHERBE, H., the effect of lipoid solvents on the rate of elimination and the carcinogenic potency of 3:4-benzpyrene after subcutaneous injection in mice, 410
——— the elimination and carcinogenic potency of 3:4-benzpyrene in mice after subcutaneous injection of non-lipoid solutions, 423